

Dynamics and distribution of doped holes in the CuO₂ plane of slightly doped antiferromagnetic YBa₂(Cu_{1-z}Li_z)₃O_{6+x} (x < 0.1) studied by Cu(1) NQR

Savinkov A., Dooglav A., Alloul H., Mendels P., Bobroff J., Collin G., Blanchard N.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Incidence of doped holes in the CuO₂ plane on the AF state was studied by Cu(1) nuclear quadrupole resonance (NQR) in slightly doped YBa₂(Cu_{1-z}Li_z)₃O_{6+x} compounds. Inhomogeneous distribution of doped holes in the plane was detected in the low temperature measurements of transverse (1/T₂) and longitudinal (1/T₁) relaxation rates. We establish that at lower T the holes motion slows down and we estimate that the holes localize finally in restricted regions (~3 lattice constants) in the Coulomb potential of the Li⁺ ions. Also we compared the hole behavior in slightly doped YBa₂(Cu_{1-z}Li_z)₃O_{6+x} samples with that in slightly doped Y_{1-y}CayBa₂Cu₃O₆. A stronger trapping potential of the in-plane Li⁺ impurities was concluded as compared to slightly doped Y_{1-y}CayBa₂Cu₃O₆ compound with out-of-plane Ca²⁺ impurities. © Pleiades Publishing, Inc., 2010.

<http://dx.doi.org/10.1134/S0021364010020074>
